

# Radial geodesics as a microscopic origin of black hole entropy.

## I: Confined under the Schwarzschild horizon

V.V.Kiselev\*†

\* Russian State Research Center “Institute for High Energy Physics”, Pobeda 1, Protvino, Moscow Region, 142281, Russia

† Moscow Institute of Physics and Technology, Institutskii per. 9, Dolgoprudnyi Moscow Region, 141700, Russia

E-mail: [kiselev@th1.ihep.su](mailto:kiselev@th1.ihep.su)

**Abstract.** Causal radial geodesics with a positive interval in the Schwarzschild metric include a subset of trajectories completely confined under a horizon, which compose a thermal statistical ensemble with the Hawking-Gibbons temperature. The Bekenstein-Hawking entropy is given by an action at corresponding geodesics of particles with a summed mass equal to that of black hole in the limit of large mass.

PACS numbers: 04.70.Dy

### 1. Introduction

At present we can observe many faces of black hole entropy first motivated by Bekenstein [1] and Hawking [2]. In addition to the thermodynamical determination by an information loss and radiation of black body, it was calculated through an action of gravitational field itself [3–5]. In a sense, a dual representation in terms of quantum fields seized by a black hole, was introduced in [6]. The most impressive evaluations were obtained in conformal field theories [7, 8] and in superstrings [9], wherein the entropy was derived by counting microstates. Other aspects were revealed by Loop Quantum Gravity [10]. Recent reviews on the subject, containing a complete list of references, can be found in [11–14].

In present paper, a new insight is offered: we maximally extend a physical space of causal geodesics of Schwarzschild solution. A notion of ‘causal’ implies a time-like value of interval on the geodesics. Then we explore the action on geodesics completely confined under a horizon in order to calculate a partition function, which actually results in the Bekenstein-Hawking relation between the area of horizon with the entropy for a particular case of Schwarzschild solution. A thermal equilibrium of confined geodesics requires a thermal quantization of levels. The space-time of geodesics confined under

the horizon is conic [15]. Thus, we present a primitive microscopic picture for an internal structure of black hole.

In section 2 we classify causal radial geodesics and thermally quantize those of confined under the horizon. Section 3 is devoted to the evaluation of entropy, while discussion, short comparison with other approaches and conclusion are collected in section 4.

## 2. Radial geodesics

In the Schwarzschild metric

$$ds^2 = \left(1 - \frac{r_g}{r}\right) dt^2 - \frac{1}{1 - \frac{r_g}{r}} dr^2 - r^2 [d\theta^2 + \sin^2 \theta d\phi^2] \quad (1)$$

the radial geodesics depend on a single integral of motion  $A$  found by Hilbert, so that

$$v^2 \equiv \left(\frac{dr}{dt}\right)^2 = \left\{1 - A \left(1 - \frac{r_g}{r}\right)\right\} \left(1 - \frac{r_g}{r}\right)^2, \quad (2)$$

where  $r_g = 2MG/c^2$  is the Schwarzschild radius, giving the well known horizon.

Indeed, a particle motion in the metric of (1) is determined by the Hamilton–Jacobi equation

$$g^{\mu\nu} \partial_\mu S_{HJ} \partial_\nu S_{HJ} - m^2 = 0, \quad (3)$$

where  $m$  denotes the particle mass. Following the general framework, we write down the solution in the form, which incorporates the single integral of the radial motion in the spherically symmetric static gravitational field,

$$S_{HJ} = -\mathcal{E} t + \mathcal{S}_{HJ}(r), \quad (4)$$

where  $\mathcal{E}$  is the conserved energy. Then, from (3) we deduce

$$\left(\frac{\partial \mathcal{S}_{HJ}}{\partial r}\right)^2 = \frac{1}{g_{tt}^2(r)} \mathcal{E}^2 - \frac{m^2}{g_{tt}(r)}, \quad \text{at} \quad g_{tt}(r) = 1 - \frac{r_g}{r}, \quad (5)$$

which results in

$$\mathcal{S}_{HJ}(r) = \int_{r_0}^{r(t)} dr \frac{1}{g_{tt}(r)} \sqrt{\mathcal{E}^2 - V^2(r)}, \quad (6)$$

where  $V^2$  is an analogue of potential,

$$V^2(r) = g_{tt}(r) m^2.$$

The trajectory is implicitly determined by equation

$$\frac{\partial S_{HJ}}{\partial \mathcal{E}} = \text{const} = -t + \int_{r_0}^{r(t)} dr \frac{1}{g_{tt}(r)} \frac{\mathcal{E}}{\sqrt{\mathcal{E}^2 - V^2(r)}}. \quad (7)$$

Taking the derivative of (7) with respect to the time<sup>‡</sup>, we get

$$1 = \dot{r} \frac{\mathcal{E}}{g_{tt}(r) \sqrt{\mathcal{E}^2 - V^2(r)}}, \quad (8)$$

resulting in (2) at  $A = m^2/\mathcal{E}^2$ .

At the trajectory the interval takes the following form:

$$ds^2 = \frac{A}{1 - A + A \frac{r_g}{r}} dr^2, \quad (9)$$

so that introducing

$$r_c = -\frac{A}{1 - A} r_g, \quad (10)$$

we get

$$ds^2 = \frac{r_c}{r_g} \frac{r}{r_c - r} dr^2. \quad (11)$$

In addition to light cones, causal geodesics are given by trajectories with a positive interval

$$ds^2 > 0 \quad \Leftrightarrow \quad \{r < r_c, r_c > 0\} \cup \{r_c < 0\}. \quad (12)$$

We subdivide the region of admissible  $r_c$ -values to three ranges or classes,

$$\begin{aligned} CI: \quad r_c < 0 & \quad \Leftrightarrow \quad 0 < A \leq 1, \\ CII: \quad r_g < r_c < +\infty & \quad \Leftrightarrow \quad 1 < A < +\infty, \\ CIII: \quad 0 < r_c < r_g & \quad \Leftrightarrow \quad -\infty < A < 0. \end{aligned} \quad (13)$$

At all of above trajectories a proper time is finite at a path under the Schwarzschild horizon, i.e. at  $r < r_g$ .

In range *CI*, the probe particle can reach a spatial infinity  $r \rightarrow +\infty$ , where its velocity is given by  $v_\infty^2 = 1 - A$ . In *CII* the particle *always* has a path outside the horizon, but the motion is restricted by  $r < r_c$  at  $r_c > r_g$ . Finally, in *CIII* the geodesics are *completely confined* under the horizon. Note, that  $r_c = r_g$  is given by  $A = \pm\infty$ , which, hence, can be identified, so that all ranges are continuously connected. This fact implies that external forces (for instance, collisions of particles) changing the gravitational integral of motion  $A$  (or  $r_c$ ), can transport or convert a trajectory from one range to another<sup>§</sup>.

The range *CII* can be completely described in isotropic Kruskal coordinates  $\{\bar{u}, \bar{v}\}$  defined by

$$\begin{cases} u = t - r_*, \\ v = t + r_*, \end{cases} \quad r_* = r + r_g \ln \left[ \frac{r}{r_g} - 1 \right], \quad (14)$$

<sup>‡</sup> As usual  $\partial_t f(t) = \dot{f}$ .

<sup>§</sup> In the quantum mechanics, the interaction should have matrix elements non-diagonal in  $A$ .

and

$$\begin{cases} \bar{u} = -2r_g e^{-\frac{u}{2r_g}}, \\ \bar{v} = +2r_g e^{+\frac{v}{2r_g}}, \end{cases} \quad (15)$$

wherein the metric takes the form

$$ds^2 = \frac{r_g}{r} \cdot e^{-\frac{r}{r_g}} \cdot d\bar{u} d\bar{v} - r^2 (d\theta^2 + \sin^2 \theta d\phi^2), \quad (16)$$

so that the singularity at the horizon  $r = r_g$  is absent, but it actually appears at  $r = 0$  as well as  $r = \infty$ , where the isotropic metric coefficient respectively becomes equal to infinity and zero (why  $CI$  cannot be completely included in the Kruskal coordinates).

The inverse transition is given by

$$\begin{cases} t = r_g \ln \left[ -\frac{\bar{v}}{\bar{u}} \right], \\ r_* = r_g \ln \left[ -\frac{\bar{u} \cdot \bar{v}}{4r_g^2} \right], \end{cases} \quad (17)$$

so that the singular point is

$$r = 0 \quad \Leftrightarrow \quad \frac{\bar{u} \cdot \bar{v}}{4r_g^2} = 1, \quad (18)$$

while the horizon corresponds to axes of  $\{\bar{u}, \bar{v}\}$ :

$$r = r_g \quad \Leftrightarrow \quad \bar{v} = 0 \cup \bar{u} = 0. \quad (19)$$

The differentials are given by

$$\begin{cases} dt = r_g \frac{d\bar{v}}{\bar{v}} - r_g \frac{d\bar{u}}{\bar{u}}, \\ dr_* = r_g \frac{d\bar{v}}{\bar{v}} + r_g \frac{d\bar{u}}{\bar{u}}, \end{cases} \quad dr_* = \frac{dr}{1 - \frac{r_g}{r}}, \quad (20)$$

which have a singularity at the horizon.

Light cones for the radial motion in metric (16) are determined by

$$d\bar{u} = 0 \quad \cup \quad d\bar{v} = 0. \quad (21)$$


The causal structure of space-time in the Kruskal coordinates is shown in Fig. 1, where a characteristic example for geodesics of  $CII$  is also depicted by a dashed curve, while a short-dashed curve represents a part of geodesic from the family of  $CI$ .

At  $A > 0$  we can easily integrate out (2) at  $r \rightarrow r_g$  to get a dependence of radial coordinate on the time near the horizon, so that

$$t - \xi r_* \rightarrow \text{const.} \quad \text{at} \quad r \rightarrow r_g, \quad (22)$$

where  $\xi = \text{sign}[\dot{r}]$  is a velocity signature. Therefore, a body falling to the horizon ( $\xi = -1$ ) has a finite value of  $\bar{v}$ , while a body coming out of the horizon ( $\xi = +1$ ) has a finite value of  $\bar{u}$  as shown in Fig.1. Further, differentials (20) give

$$\frac{d\bar{u}}{d\bar{v}} = -\frac{\bar{u}}{\bar{v}} \frac{1 - \dot{r}_*}{1 + \dot{r}_*}, \quad (23)$$



**Figure 1.** Curves of singularity  $r = 0$  restrict the maximal admissible physical region in the Kruskal variables  $\{\bar{u}, \bar{v}\}$ . The region of external observer in the Schwarzschild metric I is limited by the horizon of future  $H_+$  and of past  $H_-$ : for such the observer the future of events in the “black hole” III and past of events in the “white hole” IV are not available. An observer in region II is not causally related with an observer in I, and regions IV and III represent “black and white holes” for him respectively (in II the time  $t$  ‘flows from top to bottom’). A dashed curve is geodesic of  $CII$ : any two points at top and bottom curves of singularity can be connected by a geodesic curve in  $CII$ . A short-dashed curve shows a part of geodesic from the family of  $CI$ .

so that in vicinity of singularity

$$\frac{d\bar{u}}{d\bar{v}} \approx \frac{\bar{u}}{\bar{v}} \left( 1 - \xi \sqrt{\frac{r}{Ar_g}} \right), \quad r \rightarrow 0, \quad (24)$$


and it has the following limit independent of  $A > 0$ :

$$\frac{d\bar{u}}{d\bar{v}} = \frac{4r_g^2}{\bar{v}^2}, \quad r = 0. \quad (25)$$

These characteristics have been taken into account in Fig. 1. Note, that the isotropic Kruskal coordinates are not complete, since the trajectories coming to infinity should be described in Schwarzschild coordinates at  $r > R_0 > r_g$  (otherwise the nullification of metric coefficient will appear as an asymptotic reaching the light cone by the trajectory||).

|| An ordinary compactifying transformation can help, see [16].

Class *CIII* has a common element with class *CII*: the geodesic lines  $\bar{u} = \varkappa^2 \bar{v}$ ,  $0 < \varkappa^2 < +\infty$ . An example of common geodesics is depicted in Fig. 2. Note, that by (25) all geodesics in *CI* and *CII*, crossing the singularity at the same point of  $\{\bar{u}, \bar{v}\}$  plane, have the same tangent line, which is the extremal geodesic line at  $A \rightarrow \infty$ , as shown in Fig. 2.



**Figure 2.** A common geodesic for *CII* and *CIII*. Labels of axes are the same as in Fig. 1.

However, the space-time of geodesics in the range *CIII* cannot be described in terms of real  $\{t, r\}$  or  $\{\bar{u}, \bar{v}\}$ , since a positive interval at geodesics is expressed as

$$ds^2 = A \left(1 - \frac{r_g}{r}\right)^2 dt^2 > 0,$$

and at  $A < 0$  it gives  $dt^2 < 0$ , i.e. the imaginary differential of time.

Since we deal with causal solutions of Einstein equations for the metric with the given asymptotic behavior at  $r \rightarrow \infty$ , we have to appropriately adjust a coordinate system suitable for the description of solutions. In general, there are two options: *i*) solution should be constrained to avoid problems with some choices of coordinates, i.e. the continuation of solution is forbidden; *ii*) the solution is continuable, while coordinate systems have constrained regions of applicability. We follow the second option, which composes the paradigm of general relativity: the gravitational dynamics determines the space-time structure, i.e. causal geodesics compose the physical space-time¶.

¶ We have to specially emphasize that *i*) fixing the metric signature by  $\{+, -, -, -\}$  at any point of physical space-time as a postulate of general relativity forbids the class *CIII* of geodesics inside the black hole, *ii*) excluding the singularities on global maps (by contrast with local ones) forbids the crossing of horizons, i.e., for instance, it restricts the Schwarzschild solution by  $r > r_g$ , and the continuation of solution as the black hole is forbidden, too. Requiring the absence of singularities on a global coordinate map of space-time (in a global reference frame) is essentially equivalent to the postulate on the identity of general relativity and theory of tensor field in the Minkowski space-time. The introduction of singularities (horizons) in the global sense, i.e. the description of space-time

I shall postulate, that a full physical space-time of test particles is constructed by the following procedure of thought experiment: Imagine that a test particle is a point-like spacecraft equipped by an engine. By definition, the engine can change the integral of motion (continuously, if the system is classical, or discretely, if the system is quantum), by preserving the causality: the interval of massive particle has to be positive under the action of engine<sup>+</sup>. Thus, the space-time is what the test particle can permanently visit due to the work of engine\*. An additional constraint is an asymptotic boundary condition: a travel is starting from a region of external observer well defined by the Schwarzschild solution at large  $r > r_g$ .

So, geodesics completely confined under the horizon can be described in terms of variables  $\{\rho, \varphi_E\}$  introduced by

$$\begin{cases} \bar{u} = \varkappa i \rho e^{i\varphi_E}, \\ \bar{v} = -\frac{i}{\varkappa} \rho e^{-i\varphi_E}, \end{cases} \quad (26)$$

or (at an ‘initial’ time  $\Delta t_0 = -2r_g \ln \varkappa$ , which is set to zero)

$$\begin{cases} t = r_g \ln e^{-2i\varphi_E} = -i 2r_g \varphi_E, & \varphi_E \in [0, 2\pi], \\ r_* = 2r_g \ln \left[ -\frac{\rho}{2r_g} \right], & \rho \in [0, 2r_g], \end{cases} \quad (27)$$

so that the interval takes the form

$$ds^2 = \frac{r_g}{r} \cdot e^{-\frac{r}{r_g}} \cdot (d\rho^2 + \rho^2 d\varphi_E^2) - r^2 (d\theta^2 + \sin^2 \theta d\phi^2). \quad (28)$$

Therefore, transition (26) is periodic in  $\varphi_E$  with the period  $\Delta\varphi_E = 2\pi$ . Then, we get the period in imaginary time

$$\beta = 4\pi r_g. \quad (29)$$

We have to emphasize that transformation (27) is not a formal Wick rotation [17], since it corresponds to the space-time composed by causal geodesics continuously connected with causal geodesics in Kruskal coordinates. Thus, from (26), (27) and (29) we deduce by regular local maps, means, as we see, the refusal of the field-theoretical interpretation of general relativity and recognizing its pure geometric nature (with following problems of introducing an  $S$ -matrix in the quantum theory and so on). In the latter case we have to consider the signature of metric, which is asymptotically given as a boundary condition for the geometric equations of gravity, as the dynamical geometric quantity. Then the class of geodesics *CIII* under study is not only admissible, but it is conceptually necessary for the consistency and completeness of theory. Thus, in addition to the mathematical formalism of general relativity everybody should decide to himself (herself) which dynamical principles are admissible, acceptable or not.


<sup>+</sup> A mathematical question is a somehow smoothness or topology. I will leave it free, since a ball situated on a plane has a single common point with the plane, not a two-dimensional transition, so that a connection is not smooth nor regular, and it is topologically peculiar. Analogously, a cone situated on a plane also has a common one-dimensional line, only, and a conjugation region is not maximally smooth.

\* I do not see any ‘no-go’ principle for the existence of such the engine.

that geodesics confined under the horizon should be periodic in  $\varphi_E$  or imaginary time, and they compose a statistical ensemble with the Hawking–Gibbons temperature

$$T = \frac{1}{4\pi r_g}. \quad (30)$$

The trajectory  $\varphi_E = \pi/2$  reproduces  $\bar{u} = \varkappa^2 \bar{v}$ . Generically, the radial geodesics in range *CIII* propagate in the conic geometry of (28) (see Fig. 3), whereas  $\rho = 0 \Leftrightarrow r = r_g$ , and  $\rho = 2r_g \Leftrightarrow r = 0$ .



**Figure 3.** Physical space-times for geodesics of classes *CII* (Kruskal) and *CIII* (conic) with the common element  $\bar{u} = \bar{v}$ .

For a complete description we deduce

$$\frac{d\varphi_E}{dr} = \frac{1}{2r_g} \sqrt{1 - \frac{r_c}{r_g}} \frac{r}{r_g - r} \sqrt{\frac{r}{r_c - r}}, \quad (31)$$

which allows us to calculate a cycle change of  $\varphi_E$  with  $r$  varying from 0 to  $r_c$  and back:

$$\Delta_c \varphi_E = 2 \int_0^{r_c} dr \frac{d\varphi_E}{dr} = \frac{\pi}{2} \left[ 2 - (2 + x) \sqrt{1 - x} \right], \quad (32)$$

where we have introduced  $x = r_c/r_g$ . The ‘winding’ number (or occupation number) is defined by a number of cycles in the period, so that

$$n = \frac{2\pi}{\Delta_c \varphi_E}. \quad (33)$$

Since the motion should be periodic, we get the ‘quantization’


$$n \in \mathbb{N}. \quad (34)$$

Then, it results in the implicit quantization of  $r_c$ , i.e.  $x = x_n$ . The occupation number monotonically grows with decreasing of  $x \in [0, 1]$ . The minimal  $n$  is given by the limit  $x \rightarrow 1$ , so that from (32), (33) we deduce


$$n_{\min} = n_{x \rightarrow 1} = 2. \quad (35)$$



A picture of quantized geodesics is shown in Fig. 4.



**Figure 4.** The geodesics of *CIII* with winding numbers 2, 3, 4 in polar coordinates  $\{\rho, \varphi_E\}$ .



**Figure 5.** The quantum action.

The action of relativistic particle

$$S = -mc \int ds$$

at the geodesics confined under the horizon, is also quantized

$$S_n = -mc \cdot n \int_{\text{cycle}} dr \frac{ds}{dr} = -mc \cdot \pi r_g n \sqrt{x_n^3}, \quad (36)$$


as shown in Fig. 5.

The difference between two initial quantum actions grows with  $M$

$$S_2 - S_3 \approx \frac{\pi}{c} GMm(3 \cdot 0.924 - 2) \sim |S_2| \gg 1, \quad M \rightarrow \infty,$$

at any nonzero mass of probe particle.

Next, in addition to quantization, class *CIII* does not include light cones (excluding a consideration of non-radial motion)<sup>‡</sup>. This fact helps us to imagine the  $\{\rho, \varphi_E\}$  solid circle as ‘a compactification’ of holes in the  $\{\bar{u}, \bar{v}\}$  Kruskal coordinates: one should glue the horizons and identify them to a single point as shown in Fig. 6.



**Figure 6.** Compactifying the hole.

Finally, we have given the complete classification of radial geodesics for the Schwarzschild black hole.

### 3. Entropy

As we have just seen, the geodesics confined under the horizon periodically evolve with imaginary time. Therefore, they compose a statistical ensemble with the period equal to the inverse temperature  $\beta$  (29). The partition function of statistical ensemble is calculated as the evolution operator summed over all trajectories with the given period. For ‘classical’ particles, the classical trajectories are taken into account, only. For real time, the evolution operator is given by

$$U = \prod \sum e^{iS},$$

where  $S$  denotes the particle action, while the sum is running over the particle trajectories, and the product is taken over particles. The motion in imaginary time  $t_E = it$  gives

$$iS = i \int dt L = \int dt_E L = S_E,$$

where  $L$  is a lagrangian, and  $S_E$  is the ‘euclidian’ action. Then, the partition function is determined by

$$Z = \prod \sum e^{S_E}. \quad (37)$$

For the statistical ensemble of particles moving on geodesics confined under the horizon, the euclidian action under study is exactly the action we have just calculated.

For a given particle, as we have noted in the previous section, in the limit of large black hole mass the splitting between the action of winding number 2 and actions of

<sup>‡</sup> In fact, the light on the horizon or, more generally, massless particles can be prescribed to a trajectory being a circle around the point  $r = r_g$  with an infinitely small radius. Then, the winding number  $n = 1$  corresponds to the massless modes. We put the classical action of such modes equal to zero.

higher winding numbers is large, i.e. the action gap grows with  $M$ , so that the sum over trajectories can be approximated by the leading term of  $S_2$ :

$$\sum e^{S_E} \approx e^{S_2}, \quad (38)$$

whereas corrections are exponentially suppressed. Then, the product over the particles in the partition function is reduced to the sum of leading-term actions over particles,

$$Z \approx \prod e^{S_2} = e^{\sum S_2}.$$

The sum is easily calculated in the form

$$\sum S_2 = -2\pi c r_g \sum m. \quad (39)$$

Suppose that the gravitational field of black hole is generated by the ensemble of particles completely confined under the horizon. Then, it would be natural to suggest that the sum of particle masses is equal to the mass of black hole<sup>††</sup>:

$$M = \sum m = \frac{1}{2} r_g, \quad (40)$$

in the units  $c^2 = 1$ ,  $G = 1$ ,  $\hbar = 1$ . In these units, the sum of actions is given by

$$\sum S_2 = -\pi r_g^2 = -\frac{\beta^2}{16\pi}, \quad (41)$$

and the partition function is equal to

$$Z = \exp \left[ -\frac{\beta^2}{16\pi} \right]. \quad (42)$$

Thermodynamic relations give the following:

1. the average energy equal to the black hole mass,

$$\langle E \rangle = -\frac{\partial \ln Z}{\partial \beta} = \frac{\beta}{8\pi} = \frac{r_g}{2} = M,$$

2. the entropy

$$\mathcal{S} = \beta \langle E \rangle + \ln Z = \frac{\beta^2}{16\pi} = \pi r_g^2,$$

which is the Bekenstein–Hawking relation between the entropy and the area of horizon  $\mathcal{A} = 4\pi r_g^2$ :

$$\mathcal{S} = \frac{1}{4} \mathcal{A}.$$

Thus, in the leading approximation we have calculated the partition function of static ensemble composed by particles at the geodesics completely confined under the horizon of black hole. Putting the sum of particle masses equal to the mass of black hole we find the entropy of black hole at the microscopic level.

As for corrections to the partition function, they could be twofold. First, terms by higher winding numbers  $n$  are exponentially suppressed, so that one can easily estimate a convergency of series over  $n$ . Indeed, at large  $n$  from (32), (33) and (36) we deduce

$$S_n \rightarrow -mc\pi r_g \frac{16}{3} \sqrt[4]{\frac{3n}{16}}, \quad n \rightarrow \infty, \quad (43)$$

<sup>††</sup> Actually, the expression is fixed by a further evaluation of average energy (see below).

so that terms of series *exponentially* tend to zero with  $n$ , and series is converging.

Second, a reasonable source of corrections to the partition function is a non-radial motion, which we have not considered at all, since we believe that for the black hole of zero angular momentum the contribution of non-radial geodesics should be suppressed.

#### 4. Discussion and conclusion

We have calculated the entropy of Schwarzschild black hole by evaluating the action for radial geodesics completely confined under the horizon, which form the statistical ensemble with the Hawking temperature, in the leading order at large masses of black holes. This procedure has become reasonable under the specific quantization of trajectories by the winding number in the periodic compact variable of conic geometry induced by the confined geodesics. In this respect the Hamilton–Jacobi equation of (5) can be represented in the form

$$\frac{1}{m^2} \left( \frac{\partial \mathcal{S}_{HJ}}{\partial r_*} \right)^2 = \mathcal{E}_A - U(r), \quad \mathcal{E}_A = \frac{1}{A}, \quad U(r) = g_{tt}(r) = 1 - \frac{r_g}{r}, \quad (44)$$

which determines the spectrum of the problem, shown in Fig. 7, where the region of  $\mathcal{E}_A < 0$  represents a quantum thermal bath. Note, that all classical geodesics with  $\mathcal{E}_A < 0$  are permitted, however, a thermal equilibrium takes place if only the trajectories are periodic in the euclidian time.

The rule of quantization

$$\oint_{\Delta_c} d\varphi_E = n \cdot \int_{\Delta_c} d\varphi_E = 2\pi \quad (45)$$

can be simply rewritten in the form

$$\oint_{\Delta_c} T dt_E = n \cdot \int_{\Delta_c} T dt_E = \hbar, \quad (46)$$

where  $n = 2, 3, \dots$  is the winding number,  $T$  is the temperature, and  $\Delta_c$  denotes the contribution per cycle. One can represent (46) as


$$\oint T_n dt_E = n \hbar, \quad \Rightarrow \quad \oint dT_n dt_E = dn \cdot \hbar, \quad (47)$$

where  $T_n = nT$  is ‘the temperature associated to the  $n$ -th level of winding number’. This rule can be compared with the quasi-classical quantization by Bohr–Sommerfeld

$$\oint p dq = 2\pi n \hbar, \quad (48)$$

where  $p$  is canonically conjugated to a dynamical variable  $q$ , periodic in time.

Let us use the quasi-classics in order to count a number of microstates  $\mathcal{N}$  and to compare it with the entropy evaluated by the partition function. So, in the case under study, a role of evolution parameter is playing by the radius  $r$  (or  $\rho$ ), while the euclidian time  $t_E$  (or  $\varphi_E$ ) is the dynamical variable. It is simply to recognize that the corresponding conjugated momentum is the euclidian energy  $\mathcal{E}_n = m_n / \sqrt{-A_n}$ , where we



**Figure 7.** The gravitational potential  $U(r)$  and energy levels of particle confined under the horizon with various winding numbers.

have put index  $n$  to the particle mass to show the following: one could adiabatically vary the masses of particles composing the thermal equilibrium, when they occupy different levels with no change of entropy<sup>‡</sup>: for instance, if all particles are situated at the  $n$ -th level, the calculated product of interval per cycle  $\Delta_c s_n$  by the sum of masses at  $n$ -th level  $\sum m_n$  and winding number should be invariant,

$$n \cdot \Delta_c s_n \sum m_n c = 2 \cdot \Delta_c s_2 \sum m_2 c \equiv -S_2.$$

The invariance of action under such the re-distributions of particles between the levels conserves the entropy. This variation is responsible for the entropy as a measure of chaos. Then, using the definition of temperature as an average energy per particle

$$T_n = \frac{E_n}{\nu}, \quad E_n = \sum \mathcal{E}_n,$$

<sup>‡</sup> The action of particle can be rewritten in the form  $S = -mc \int dr L_r$ , at  $L_r = \sqrt{g_{\mu\nu} dx^\mu / dr dx^\nu / dr}$ , so that the momentum conjugated to  $t_E$  is  $p_E = -mc \partial L_r / \partial (dt_E / dr)$ , which is equal to  $E$ .

where  $\nu$  is a number of particles at the  $n$ -th level, ‘in a quasi-classical manner’ of (47) by (48) we get

$$\frac{1}{\nu} \oint E_n dt_E = \hbar n, \quad dE_n \oint dt_E = \hbar \nu dn, \quad (49)$$

Thermodynamically, we write down

$$dE_n = T_n dS = n T dS$$

and deduce

$$T dS \oint dt_E = \hbar \nu \frac{dn}{n} = \hbar d \ln \mathcal{N}, \quad (50)$$

where  $\mathcal{N} = n^\nu$  is the total number of microstates for  $\nu$  particles each having  $n$  states. Integrating (50) over all levels, particles and their masses, we obtain

$$\int T dS \oint dt_E = \hbar \int_2^{\sqrt[\nu]{\mathcal{N}}} \nu \frac{dn}{n} \approx \hbar \ln \mathcal{N}, \quad \mathcal{N}^{1/\nu} \gg 1. \quad (51)$$

On the other hand, the quantity

$$\mathcal{I} = \int T dS \oint dt_E = \hbar \int dS = \hbar \mathcal{S} \quad (52)$$

is an adiabatic invariant value independent of  $n$ , i.e. the entropy multiplied by  $\hbar$ .

Our calculations have given

$$\mathcal{I} = \hbar \frac{\pi r_g^2}{\ell_{\text{Pl}}^2}, \quad (53)$$

where we have restored the fundamental units and denoted the square of Planck length by  $\ell_{\text{Pl}}^2 = G\hbar/c^3$ . Comparison of (52) with (51) gives the standard correspondence between the entropy and number of microstates,

$$\mathcal{S} = \ln \mathcal{N}. \quad (54)$$

Of course, we may treat the logical chain (49)–(54) backward, to derive the quantization rule (49) from the known expression for the entropy.

The winding number is restricted by a maximal value, since  $\nu \geq 1$ , and hence, the limit is  $n_{\text{max}} = \mathcal{N} = \exp \mathcal{S} \gg 1$ . Let us remind that the action gap was essential for the derivation of partition function, so that  $Mm/m_{\text{Pl}}^2 \gg 1$ , and the particle mass is restricted from bottom. At  $n \gg 1$  the appropriate mass tends to zero (see (43))§, and it should reach a minimal admissible value, so that  $\nu \gg 1$ .

Classical trajectories unsatisfying (45) are not forbidden, but they do not belong to a thermal bath with a definite temperature, i.e. they are not in the thermal equilibrium.

Such the quantization intrinsically differs from the quasi-classics in the quantum mechanics, since in nature it unavoidably involves the temperature: it seems that any real theory of quantum gravity should by origin include the temperature (or statistical

§ Note also, that at  $n \rightarrow n_{\text{max}}$  the energy of particle gets the value about the black hole mass  $E_{\mathcal{N}} = m_{\mathcal{N}}/\sqrt{-A_{\mathcal{N}}} \approx 3M/8$ .

chaos?). The exterior of black hole, i.e. the space-time outside the horizon, forms the thermostats of thermal geodesics confined under the horizon.

Having calculated the entropy by the partition function, we have neglected the contributions by higher winding numbers (or used the invariance of action by adjusting the sum of masses to get the given mass of black hole) as well as by a non-radial motion.

The leading term with the minimal winding number gives the state, which is ‘coherent’ in a sense, that the geodesic, connecting  $r_g$  to  $r = 0$ , matches all  $r$  to the point  $r = r_g$ . Due to the coherence, the physics under the horizon can be completely described by the physics on the surface of the horizon. This fact reveals the principle of holography: we do not need a three dimensional interior, its surface is sufficient [18]. The information inside the hole is equivalent to its snapshot of horizon surface. Therefore, the calculation of entropy based on the two dimensional conformal field theory (2D CFT) by the Cardy formula for the number of states [19] is not in a straightforwardly visible contradiction with the given representation. Though we cannot find a direct connection between our approach and the method based on 2D CFT, since we have not introduced quantum fields. The same note concerns for the comparison with the superstring calculation of entropy by counting appropriate microstates (the vehicle is armed by the Cardy formula, again).

The construction offered is, in a sense, dual to the Hawking calculation based on the evaluation of gravitational action for the given metric itself, since we deal with the action for the particles moving in the metric. Therefore, we have implicitly supposed that the metric is an effective average collective gravitational field of particles. A duality is also suggestive in connection to a method by Frolov and Novikov, who constructed a calculation separating visible and invisible modes of quantum field [6]. The former is radiated by the black hole, while the later is eaten by the hole. In this sense, the entropy is given by the information eaten by the black hole.

In conclusion, we have presented a picture for the interior of Schwarzschild black hole, that composes the first part of serial article, which will be continued by an analogous consideration of Reissner–Nordström black hole.

The author expresses a gratitude to academician A.A.Logunov, who has asked me for a description of radial geodesics in the metric of Schwarzschild black hole. I specially thank associated professor O.I.Tolstikhin for a useful remark.

This work is partially supported by the grant of the president of Russian Federation for scientific schools NSc-1303.2003.2, and the Russian Foundation for Basic Research, grants 04-02-17530, 05-02-16315.

## References

- [1] J. D. Bekenstein, Lett. Nuovo Cim. **4** (1972) 737;  
J. D. Bekenstein, Phys. Rev. D **7** (1973) 2333;  
J. D. Bekenstein, Phys. Rev. D **9** (1974) 3292.
- [2] S. W. Hawking, Commun. Math. Phys. **43** (1975) 199.

- [3] S. W. Hawking, Phys. Rev. D **13** (1976) 191;  
G. W. Gibbons and S. W. Hawking, Phys. Rev. D **15** (1977) 2738;  
S. W. Hawking and G. T. Horowitz, Class. Quant. Grav. **13** (1996) 1487 [arXiv:gr-qc/9501014].
- [4] S. Carlip, Phys. Rev. Lett. **82** (1999) 2828 [arXiv:hep-th/9812013];  
S. N. Solodukhin, Phys. Lett. B **454** (1999) 213 [arXiv:hep-th/9812056].
- [5] V. O. Solovev, Phys. Rev. D **61** (2000) 027502 [arXiv:hep-th/9905220].
- [6] V. P. Frolov and I. Novikov, Phys. Rev. D **48** (1993) 4545 [arXiv:gr-qc/9309001];  
V. P. Frolov and I. Novikov, “Black hole physics: basic concepts and new developments” —  
Dordrecht: Kluwer Academic, 1998.
- [7] S. Carlip, Phys. Rev. Lett. **88** (2002) 241301 [arXiv:gr-qc/0203001];  
M. I. Park, Nucl. Phys. B **634** (2002) 339 [arXiv:hep-th/0111224].
- [8] A. Strominger, JHEP **9802** (1998) 009 [arXiv:hep-th/9712251].
- [9] A. Strominger and C. Vafa, Phys. Lett. B **379** (1996) 99 [arXiv:hep-th/9601029];  
J. M. Maldacena and A. Strominger, Phys. Rev. Lett. **77** (1996) 428 [arXiv:hep-th/9603060];  
G. T. Horowitz and A. Strominger, Phys. Rev. Lett. **77** (1996) 2368 [arXiv:hep-th/9602051].  
C. G. Callan and J. M. Maldacena, Nucl. Phys. B **472** (1996) 591 [arXiv:hep-th/9602043];  
R. Dijkgraaf, E. Verlinde and H. Verlinde, Nucl. Phys. B **486** (1997) 77 [arXiv:hep-th/9603126];  
C. V. Johnson, R. R. Khuri and R. C. Myers, Phys. Lett. B **378** (1996) 78 [arXiv:hep-th/9603061];  
J. M. Maldacena, A. Strominger and E. Witten, JHEP **9712** (1997) 002 [arXiv:hep-th/9711053].
- [10] A. Ashtekar, J. Baez, A. Corichi and K. Krasnov, Phys. Rev. Lett. **80**, 904 (1998)  
[arXiv:gr-qc/9710007].
- [11] R. Brout, S. Massar, R. Parentani and P. Spindel, Phys. Rept. **260** (1995) 329.
- [12] T. Damour, arXiv:hep-th/0401160.
- [13] D. V. Fursaev, arXiv:gr-qc/0404038.
- [14] S. Das, arXiv:hep-th/0403202.
- [15] L. Susskind, arXiv:hep-th/9309145;  
S. Carlip and C. Teitelboim, Class. Quant. Grav. **12** (1995) 1699 [arXiv:gr-qc/9312002];  
C. Callan and F. Wilczek, Phys. Lett. B **333** (1994) 55 [arXiv:hep-th/9401072].
- [16] N.D.Birrell, P.C.W.Davies, “Quantum fields in curved space” – Cambridge: Cambridge University  
Press, 1982.
- [17] J. W. York, Phys. Rev. D **33** (1986) 2092.
- [18] L. Susskind, J. Math. Phys. **36**, 6377 (1995) [arXiv:hep-th/9409089];  
R. Bousso, Rev. Mod. Phys. **74** (2002) 825 [arXiv:hep-th/0203101].
- [19] J. L. Cardy, Nucl. Phys. B **270** (1986) 186;  
H. W. J. Bloete, J. L. Cardy and M. P. Nightingale, Phys. Rev. Lett. **56** (1986) 742.